

# The Riemann-Roch Theorem

TIFR  
Mumbai, India

Paul Baum  
Penn State

7 August, 2015

Five lectures:

1. Dirac operator✓
2. Atiyah-Singer revisited✓
3. What is K-homology?✓
4. Beyond ellipticity✓
5. The Riemann-Roch theorem

## THE RIEMANN-ROCH THEOREM

1. Classical Riemann-Roch ✓
2. Hirzebruch-Riemann-Roch (HRR) ✓
3. Grothendieck-Riemann-Roch (GRR)
4. RR for possibly singular varieties (Baum-Fulton-MacPherson)

## REFERENCES

P. Baum, W. Fulton, and R. MacPherson *Riemann-Roch for singular varieties* Publ. Math. IHES 45: 101-167, 1975.

P. Baum, W. Fulton, and R. MacPherson *Riemann-Roch and topological K-theory for singular varieties* Acta Math. 143: 155-192, 1979.

P. Baum, W. Fulton, and G. Quart *Lefschetz-Riemann-Roch for singular varieties* Acta Math. 143: 193-211, 1979.

## HIRZEBRUCH-RIEMANN-ROCH

$M$  non-singular projective algebraic variety /  $\mathbb{C}$

$E$  an algebraic vector bundle on  $M$

$\underline{E}$  = sheaf of germs of algebraic sections of  $E$

$H^j(M, \underline{E}) := j$ -th cohomology of  $M$  using  $\underline{E}$ ,  
 $j = 0, 1, 2, 3, \dots$

### LEMMA

For all  $j = 0, 1, 2, \dots$   $\dim_{\mathbb{C}} H^j(M, \underline{E}) < \infty$ .

For all  $j > \dim_{\mathbb{C}}(M)$ ,  $H^j(M, \underline{E}) = 0$ .

---

$$\chi(M, E) := \sum_{j=0}^n (-1)^j \dim_{\mathbb{C}} H^j(M, \underline{E})$$

$$n = \dim_{\mathbb{C}}(M)$$

---

THEOREM[HRR] Let  $M$  be a non-singular projective algebraic variety /  $\mathbb{C}$  and let  $E$  be an algebraic vector bundle on  $M$ . Then

$$\chi(M, E) = (ch(E) \cup Td(M))[M]$$

# Hirzebruch-Riemann-Roch

## Theorem (HRR)

*Let  $M$  be a non-singular projective algebraic variety /  $\mathbb{C}$  and let  $E$  be an algebraic vector bundle on  $M$ . Then*

$$\chi(M, E) = (ch(E) \cup Td(M))[M]$$

EXAMPLE. Let  $M$  be a compact complex-analytic manifold.

Set  $\Omega^{p,q} = C^\infty(M, \Lambda^{p,q} T^* M)$

$\Omega^{p,q}$  is the  $\mathbb{C}$  vector space of all  $C^\infty$  differential forms of type  $(p, q)$

Dolbeault complex

$$0 \longrightarrow \Omega^{0,0} \longrightarrow \Omega^{0,1} \longrightarrow \Omega^{0,2} \longrightarrow \dots \longrightarrow \Omega^{0,n} \longrightarrow 0$$

The Dirac operator (of the underlying  $\text{Spin}^c$  manifold) is the assembled Dolbeault complex

$$\bar{\partial} + \bar{\partial}^*: \bigoplus_j \Omega^{0,2j} \longrightarrow \bigoplus_j \Omega^{0,2j+1}$$

The index of this operator is the arithmetic genus of  $M$  — i.e. is the Euler number of the Dolbeault complex.



Let  $X$  be a finite CW complex.

The three versions of  $K$ -homology are isomorphic.

$$K_j^{\text{homotopy}}(X) \begin{array}{c} \xrightarrow{\quad} \\ \xleftarrow{\quad} \end{array} K_j(X) \longrightarrow KK^j(C(X), \mathbb{C})$$

homotopy theory

$K$ -cycles

Atiyah-BDF-Kasparov

$$j = 0, 1$$

Let  $X$  be a finite CW complex.

The three versions of  $K$ -homology are isomorphic.

$$K_j^{\text{homotopy}}(X) \begin{array}{c} \xrightarrow{\quad} \\ \xleftarrow{\quad} \end{array} K_j(X) \longrightarrow KK^j(C(X), \mathbb{C})$$

homotopy theory

$K$ -cycles

Atiyah-BDF-Kasparov

$$j = 0, 1$$

$X$  is a finite CW complex.

### CHERN CHARACTER

The Chern character is often viewed as a functorial map of contravariant functors :

$$ch: K^j(X) \longrightarrow \bigoplus_l H^{j+2l}(X; \mathbb{Q})$$
$$j = 0, 1$$

Note that this is a map of rings.

$X$  is a finite CW complex.

A more inclusive (and more accurate) view of the Chern character is that it is a pair of functorial maps :

$$ch: K^j(X) \longrightarrow \bigoplus_l H^{j+2l}(X; \mathbb{Q}) \quad \textit{contravariant}$$

$$ch_{\#}: K_j(X) \longrightarrow \bigoplus_l H_{j+2l}(X; \mathbb{Q}) \quad \textit{covariant}$$

$K_*(X)$  is a module over  $K^*(X)$ .

$H_*(X; \mathbb{Q})$  is a module over  $H^*(X; \mathbb{Q})$ . cap product

The Chern character respects these module structures.

Definition of the Chern character in homology  $j = 0, 1$

$$ch_{\#} : K_j(X) \longrightarrow \bigoplus_l H_{j+2l}(X; \mathbb{Q}) \text{ covariant}$$

$$ch_{\#}(M, E, \varphi) := \varphi_*(ch(E) \cup Td(M)) \cap [M]$$

$$\varphi_* : H_*(M; \mathbb{Q}) \longrightarrow H_*(X; \mathbb{Q})$$

$K_*(X)$  is a module over  $K^*(X)$ .

Let  $(M, E, \varphi)$  be a  $K$ -cycle on  $X$ .

Let  $F$  be a  $\mathbb{C}$  vector bundle on  $X$ .

Then:

$$F \cdot (M, E, \varphi) := (M, E \otimes \varphi^*(F), \varphi)$$

and the module structure is respected :

$$ch_{\#}(F \cdot (M, E, \varphi)) = ch(F) \cap ch_{\#}(M, E, \varphi)$$

## $K$ -theory and $K$ -homology in algebraic geometry

Let  $X$  be a (possibly singular) projective algebraic variety  $/\mathbb{C}$ .

Grothendieck defined two abelian groups:

$K_{alg}^0(X)$  = Grothendieck group of algebraic vector bundles on  $X$ .

$K_0^{alg}(X)$  = Grothendieck group of coherent algebraic sheaves on  $X$ .

$K_{alg}^0(X)$  = the algebraic geometry  $K$ -theory of  $X$  **contravariant**.

$K_0^{alg}(X)$  = the algebraic geometry  $K$ -homology of  $X$  **covariant**.





## $K$ -theory in algebraic geometry

$\text{Vect}_{alg}X =$

set of isomorphism classes of algebraic vector bundles on  $X$ .

$A(\text{Vect}_{alg}X) =$  free abelian group

with one generator for each element  $[E] \in \text{Vect}_{alg}X$ .

For each short exact sequence  $\xi$

$$0 \rightarrow E' \rightarrow E \rightarrow E'' \rightarrow 0$$

of algebraic vector bundles on  $X$ , let  $r(\xi) \in A(\text{Vect}_{alg}X)$  be

$$r(\xi) := [E'] + [E''] - [E]$$

## $K$ -theory in algebraic geometry

$\mathcal{R} \subset A(\text{Vect}_{alg}(X))$  is the subgroup of  $A(\text{Vect}_{alg}X)$  generated by all  $r(\xi) \in A(\text{Vect}_{alg}X)$ .

DEFINITION.  $K_{alg}^0(X) := A(\text{Vect}_{alg}X)/\mathcal{R}$

Let  $X, Y$  be (possibly singular) projective algebraic varieties  $/\mathbb{C}$ .  
Let

$$f: X \longrightarrow Y$$

be a morphism of algebraic varieties.

Then have the map of abelian groups

$$f^*: K_{alg}^0(X) \longleftarrow K_{alg}^0(Y)$$

$$[f^*E] \longleftarrow [E]$$

Vector bundles pull back.  $f^*E$  is the pull-back via  $f$  of  $E$ .

## $K$ -homology in algebraic geometry

$$\mathcal{S}_{alg}X =$$

set of isomorphism classes of coherent algebraic sheaves on  $X$ .

$$A(\mathcal{S}_{alg}X) = \text{free abelian group}$$

with one generator for each element  $[\mathcal{E}] \in \mathcal{S}_{alg}X$ .

For each short exact sequence  $\xi$

$$0 \rightarrow \mathcal{E}' \rightarrow \mathcal{E} \rightarrow \mathcal{E}'' \rightarrow 0$$

of coherent algebraic sheaves on  $X$ , let  $r(\xi) \in A(\mathcal{S}_{alg}X)$  be

$$r(\xi) := [\mathcal{E}'] + [\mathcal{E}'' ] - [\mathcal{E}]$$

## $K$ -homology in algebraic geometry

$\mathfrak{R} \subset A(\mathcal{S}_{alg}(X))$  is the subgroup of  $A(\mathcal{S}_{alg}X)$  generated by all  $r(\xi) \in A(\mathcal{S}_{alg}X)$ .

DEFINITION.  $K_0^{alg}(X) := A(\mathcal{S}_{alg}X)/\mathfrak{R}$

Let  $X, Y$  be (possibly singular) projective algebraic varieties  $/\mathbb{C}$ .

Let

$$f: X \longrightarrow Y$$

be a morphism of algebraic varieties.

Then have the map of abelian groups

$$f_*: K_0^{alg}(X) \longrightarrow K_0^{alg}(Y)$$

$$[\mathcal{E}] \mapsto \sum_j (-1)^j [(R^j f)\mathcal{E}]$$

$f: X \rightarrow Y$  morphism of algebraic varieties

$\mathcal{E}$  coherent algebraic sheaf on  $X$

For  $j \geq 0$ , define a presheaf  $(W^j f)\mathcal{E}$  on  $Y$  by

$$U \mapsto H^j(f^{-1}U; \mathcal{E}|_{f^{-1}U}) \quad U \text{ an open subset of } Y$$

Then

$$(R^j f)\mathcal{E} := \text{the sheafification of } (W^j f)\mathcal{E}$$

$f: X \rightarrow Y$  morphism of algebraic varieties

$$f_*: K_0^{alg}(X) \longrightarrow K_0^{alg}(Y)$$

$$[\mathcal{E}] \mapsto \sum_j (-1)^j [(R^j f)\mathcal{E}]$$

SPECIAL CASE of  $f_*: K_0^{alg}(X) \longrightarrow K_0^{alg}(Y)$

$Y$  is a point.  $Y = \cdot$

$\epsilon: X \rightarrow \cdot$  is the map of  $X$  to a point.

$$K_{alg}^0(\cdot) = K_0^{alg}(\cdot) = \mathbb{Z}$$

$$\epsilon_*: K_0^{alg}(X) \rightarrow K_0^{alg}(\cdot) = \mathbb{Z}$$

$$\epsilon_*(\mathcal{E}) = \chi(X; \mathcal{E}) = \sum_j (-1)^j \dim_{\mathbb{C}} H^j(X; \mathcal{E})$$

$$X \text{ non-singular} \implies K_{alg}^0(X) \cong K_0^{alg}(X)$$

Let  $X$  be non-singular.

Let  $E$  be an algebraic vector bundle on  $X$ .

$\underline{E}$  denotes the sheaf of germs of algebraic sections of  $E$ .

Then  $E \mapsto \underline{E}$  is an isomorphism of abelian groups

$$K_{alg}^0(X) \longrightarrow K_0^{alg}(X)$$

This is Poincaré duality within the context of algebraic geometry  
K-theory&K-homology.



$$X \text{ non-singular} \implies K_{alg}^0(X) \cong K_0^{alg}(X)$$

Let  $X$  be non-singular.

The inverse map

$$K_0^{alg}(X) \rightarrow K_{alg}^0(X)$$

is defined as follows.

Let  $\mathcal{F}$  be a coherent algebraic sheaf on  $X$ .

Since  $X$  is non-singular,

$\mathcal{F}$  has a finite resolution by algebraic vector bundles.

$$X \text{ non-singular} \implies K_{alg}^0(X) \cong K_0^{alg}(X)$$

$\mathcal{F}$  has a finite resolution by algebraic vector bundles.

i.e.  $\exists$  algebraic vector bundles on  $X$   $E_r, E_{r-1}, \dots, E_0$  and an exact sequence of coherent algebraic sheaves

$$0 \rightarrow \underline{E}_r \rightarrow \underline{E}_{r-1} \rightarrow \dots \rightarrow \underline{E}_0 \rightarrow \mathcal{F} \rightarrow 0$$

Then  $K_0^{alg}(X) \rightarrow K_{alg}^0(X)$  is

$$\mathcal{F} \mapsto \sum_j (-1)^j E_j$$

# Grothendieck-Riemann-Roch

## Theorem (GRR)

*Let  $X, Y$  be non-singular projective algebraic varieties  $/\mathbb{C}$ , and let  $f : X \rightarrow Y$  be a morphism of algebraic varieties. Then there is commutativity in the diagram :*

$$\begin{array}{ccc} K_{alg}^0(X) & \longrightarrow & K_{alg}^0(Y) \\ ch(\ ) \cup Td(X) & \downarrow & \downarrow ch(\ ) \cup Td(Y) \\ H^*(X; \mathbb{Q}) & \longrightarrow & H^*(Y; \mathbb{Q}) \end{array}$$

## WARNING!!!

The horizontal arrows in the GRR commutative diagram

$$\begin{array}{ccc} K_{alg}^0(X) & \longrightarrow & K_{alg}^0(Y) \\ ch() \cup Td(X) & \downarrow & \downarrow ch() \cup Td(Y) \\ H^*(X; \mathbb{Q}) & \longrightarrow & H^*(Y; \mathbb{Q}) \end{array}$$

are wrong-way (i.e. Gysin) maps.

$$K_{alg}^0(X) \cong K_0^{alg}(X) \xrightarrow{f_*} K_0^{alg}(Y) \cong K_{alg}^0(Y)$$

$$H^*(X; \mathbb{Q}) \cong H_*(X; \mathbb{Q}) \xrightarrow{f_*} H_*(Y; \mathbb{Q}) \cong H^*(Y; \mathbb{Q})$$

Poincaré duality

Poincaré duality

# Riemann-Roch for possibly singular complex projective algebraic varieties

Let  $X$  be a (possibly singular) projective algebraic variety /  $\mathbb{C}$

Then (Baum-Fulton-MacPherson) there are functorial maps

$\alpha_X : K_{alg}^0(X) \longrightarrow K_{top}^0(X)$       *K-theory*      *contravariant*  
natural transformation of contravariant functors

$\beta_X : K_0^{alg}(X) \longrightarrow K_0^{top}(X)$       *K-homology*      *covariant*  
natural transformation of covariant functors

Everything is natural. No wrong-way (i.e. Gysin) maps are used.

$$\alpha_X: K_{alg}^0(X) \longrightarrow K_{top}^0(X)$$

is the forgetful map which sends an algebraic vector bundle  $E$  to the underlying topological vector bundle of  $E$ .

$$\alpha_X(E) := E_{\text{topological}}$$

Let  $X, Y$  be projective algebraic varieties  $/\mathbb{C}$ , and let  $f : X \rightarrow Y$  be a morphism of algebraic varieties. Then there is commutativity in the diagram :

$$\begin{array}{ccc} K_{alg}^0(X) & \longleftarrow & K_{alg}^0(Y) \\ \alpha_X \downarrow & & \downarrow \alpha_Y \\ K_{top}^0(X) & \longleftarrow & K_{top}^0(Y) \end{array}$$

i.e. natural transformation of contravariant functors

Let  $X, Y$  be projective algebraic varieties  $/\mathbb{C}$ , and let  $f : X \rightarrow Y$  be a morphism of algebraic varieties. Then there is commutativity in the diagram :

$$\begin{array}{ccc} K_{alg}^0(X) & \longleftarrow & K_{alg}^0(Y) \\ \alpha_X \downarrow & & \downarrow \alpha_Y \\ K_{top}^0(X) & \longleftarrow & K_{top}^0(Y) \\ ch \downarrow & & \downarrow ch \\ H^*(X; \mathbb{Q}) & \longleftarrow & H^*(Y; \mathbb{Q}) \end{array}$$



Let  $X, Y$  be projective algebraic varieties  $/\mathbb{C}$ , and let  $f : X \rightarrow Y$  be a morphism of algebraic varieties. Then there is commutativity in the diagram :

$$\begin{array}{ccc} K_0^{alg}(X) & \longrightarrow & K_0^{alg}(Y) \\ \beta_X \downarrow & & \downarrow \beta_Y \\ K_0^{top}(X) & \longrightarrow & K_0^{top}(Y) \end{array}$$

i.e. natural transformation of covariant functors

Notation.  $K_*^{top}$  is  $K$ -cycle  $K$ -homology.

Let  $X, Y$  be projective algebraic varieties  $/\mathbb{C}$ , and let  $f : X \rightarrow Y$  be a morphism of algebraic varieties. Then there is commutativity in the diagram :

$$\begin{array}{ccc} K_{alg}^0(X) & \longleftarrow & K_{alg}^0(Y) \\ \alpha_X \downarrow & & \downarrow \alpha_Y \\ K_{top}^0(X) & \longleftarrow & K_{top}^0(Y) \\ ch \downarrow & & \downarrow ch \\ H^*(X; \mathbb{Q}) & \longleftarrow & H^*(Y; \mathbb{Q}) \end{array}$$

Let  $X, Y$  be projective algebraic varieties  $/\mathbb{C}$ , and let  $f : X \rightarrow Y$  be a morphism of algebraic varieties. Then there is commutativity in the diagram :

$$\begin{array}{ccc} K_0^{alg}(X) & \longrightarrow & K_0^{alg}(Y) \\ \beta_X \downarrow & & \downarrow \beta_Y \\ K_0^{top}(X) & \longrightarrow & K_0^{top}(Y) \\ ch_{\#} \downarrow & & \downarrow ch_{\#} \\ H_*(X; \mathbb{Q}) & \longrightarrow & H_*(Y; \mathbb{Q}) \end{array}$$

## Definition of $\beta_X: K_0^{alg}(X) \rightarrow K_0^{top}(X)$

Let  $\mathcal{F}$  be a coherent algebraic sheaf on  $X$ .

Choose an embedding of projective algebraic varieties

$$\iota: X \hookrightarrow W$$

where  $W$  is non-singular.

$\iota_*\mathcal{F}$  is the push forward (i.e. extend by zero) of  $\mathcal{F}$ .

$\iota_*\mathcal{F}$  is a coherent algebraic sheaf on  $W$ .

$\iota_*\mathcal{F}$  is a coherent algebraic sheaf on  $W$ .

Since  $W$  is non-singular,  $\iota_*\mathcal{F}$  has a finite resolution by algebraic vector bundles.

$$0 \rightarrow \underline{E}_r \rightarrow \underline{E}_{r-1} \rightarrow \dots \rightarrow \underline{E}_0 \rightarrow \iota_*\mathcal{F} \rightarrow 0$$

Consider

$$0 \rightarrow E_r \rightarrow E_{r-1} \rightarrow \dots \rightarrow E_0 \rightarrow 0$$

These are algebraic vector bundles on  $W$  and maps of algebraic vector bundles such that for each  $p \in W - \iota(X)$  the sequence of finite dimensional  $\mathbb{C}$  vector spaces

$$0 \rightarrow (E_r)_p \rightarrow (E_{r-1})_p \rightarrow \dots \rightarrow (E_0)_p \rightarrow 0$$

is exact.

Choose Hermitian structures for  $E_r, E_{r-1}, \dots, E_0$   
Then for each vector bundle map

$$\sigma: E_j \rightarrow E_{j-1}$$

there is the adjoint map

$$\sigma^*: E_j \leftarrow E_{j-1}$$

$$\sigma \oplus \sigma^*: \bigoplus_j E_{2j} \longrightarrow \bigoplus_j E_{2j+1}$$

is a map of topological vector bundles which is an isomorphism on  $W - \iota(X)$ .

Let  $\Omega$  be an open set in  $W$  with smooth boundary  $\partial\Omega$  such that  $\bar{\Omega} = \Omega \cup \partial\Omega$  is a compact manifold with boundary which retracts onto  $\iota(X)$ .  $\bar{\Omega} \rightarrow \iota(X)$ .

Set

$$M = \bar{\Omega} \cup_{\partial\Omega} \bar{\Omega}$$

$M$  is a closed  $\text{Spin}^c$  manifold which maps to  $X$  by:

$$\varphi: M = \bar{\Omega} \cup_{\partial\Omega} \bar{\Omega} \rightarrow \bar{\Omega} \rightarrow \iota(X) = X$$

On  $M = \bar{\Omega} \cup_{\partial\Omega} \bar{\Omega}$  let  $E$  be the topological vector bundle

$$E = \bigoplus_j E_{2j} \cup_{(\sigma \oplus \sigma^*)} \bigoplus_j E_{2j+1}$$

Then  $\beta_X : K_0^{alg}(X) \rightarrow K_0^{top}(X)$  is :

$$\mathcal{F} \mapsto (M, E, \varphi)$$

$$M = \bar{\Omega} \cup_{\partial\Omega} \bar{\Omega}$$



Equivalent definition of  $\beta_X: K_0^{alg}(X) \rightarrow K_0^{top}(X)$

Let  $(M, E, \varphi)$  be an algebraic  $K$ -cycle on  $X$ , i.e.

- $M$  is a non-singular complex projective algebraic variety.
- $E$  is an algebraic vector bundle on  $M$ .
- $\varphi: M \rightarrow X$  is a morphism of projective algebraic varieties.

Then:

$$\beta_X(\varphi_*(\underline{E})) = (M, E, \varphi)_{\text{topological}}$$

## Module structure

$K_{alg}^0(X)$  is a ring and  $K_0^{alg}(X)$  is a module over this ring.

$\alpha_X: K_{alg}^0(X) \rightarrow K_{top}^0(X)$  is a homomorphism of rings.

$\beta_X: K_0^{alg}(X) \rightarrow K_0^{top}(X)$  respects the module structures.

## Todd class

Set

$$\mathrm{td}(X) = \mathrm{ch}(\beta_X(\mathcal{O}_X)) \quad \mathrm{td}(X) \in H_*(X; \mathbb{Q})$$

If  $X$  is non-singular, then  $\mathrm{td}(X) = \mathrm{Todd}(X) \cap [X]$ .

With  $X$  possibly singular and  $E$  an algebraic vector bundle on  $X$

$$\chi(X, \underline{E}) = \epsilon_*(\mathrm{ch}(E) \cap \mathrm{td}(X))$$

$\epsilon: X \rightarrow \cdot$  is the map of  $X$  to a point.

$$\epsilon_*: H_*(X; \mathbb{Q}) \rightarrow H_*(\cdot; \mathbb{Q}) = \mathbb{Q}$$

Let

$$\begin{array}{c} \tilde{X} \\ \downarrow \pi \\ X \end{array}$$

be resolution of singularities in the sense of Hironaka.

$$\pi_*: H_*(\tilde{X}; \mathbb{Q}) \rightarrow H_*(X; \mathbb{Q})$$

Lemma.  $\pi_*(Td(\tilde{X}) \cap [\tilde{X}])$  is intrinsic to  $X$  i.e. does not depend on the choice of the resolution of singularities.

$td(X) \in H_*(X; \mathbb{Q})$  is also intrinsic to  $X$ .

$td(X) - \pi_*(Td(\tilde{X}) \cap [\tilde{X}])$  is given by a homology class on  $X$  which (in a canonical way) is supported on the singular locus of  $X$ .

Problem. In examples calculate  $td(X) \in H_*(X; \mathbb{Q})$ .

For toric varieties see papers of J. Shaneson and S. Cappell.